




NeoTell


1



2




3



4



5



6



7





SCIENTIFIC REFERENCES

1. Sennerby L. Implantologie 2013;21(1):21-33
2. Kokovic V, Jung R, Feloutzis A, Todorovic V, Jurisic M, 2. Härmämerle C. Clinical Oral Implants Research, 00, 2013, 1-6
3. Bonstein M, Hart C, Halbritter S, Morton D, Buser D. Clin Implant Dent Relat Res 2009
4. Serge Baltayan, Joan Pi-Anfuns, Tara Aghaloo, Peter Moy. J Oral Maxillofac Surg 74:1145-1152, 2016
5. Östman P-O. Clinical Implant Dentistry and Related. Research, Volume 7, Supplement 1, 2005
6. Rodrigo D, Aracil L, Martin C, Sanz M. Clin. Oral Impl. Res. 21, 2010; 255-261
7. Pagliani L, Sennerby L, Petersson A, Verrochi D, Volpe S & Andersson P. Journal of Oral Rehabilitation 2012
8. Trisi P, Carlesi T, Colagiovanni M, Perfetti G. Journal of Osteology and Biomaterials, Volume 1, Number 3, 2010
9. Stefan Paul Hicklin, Esther Schneebeli, Vivianne Chappuis, Simone Francesco Marco Janner, Daniel Buser, Urs Brägger. Clin. Oral Impl. Res 00, 2015; 1-9



The above is a summary of scientific data and not an official recommendation by Neoss.
To monitor osseointegration measure at placement and before final restoration.

15603_0 EN 23-04 © Neoss Group, 2023. Copyrights, design rights and trademarks, Neoss documents, software and designs may not be reprinted, copied or published in whole or part, without the written authorization of Neoss. Neoss, the N logo and NeoTell™ are trademarks of Neoss Group.